Sandblasting the r-Process: Spallation of Ejecta from Neutron Star Mergers

XILU WANG^{1, 2, *}

(N3AS COLLABORATION)

Brian D. Fields, 3,4 Matthew Mumpower, 5,6 Trevor Sprouse, 2,5 Rebecca Surman, 2 and Nicole Vassh 2

¹Department of Physics, University of California, Berkeley, CA 94720, USA
 ²Department of Physics, University of Notre Dame, Notre Dame, IN 46556, USA
 ³Department of Astronomy, University of Illinois at Urbana-Champaign, Urbana, IL 61801, USA
 ⁴Department of Physics, University of Illinois at Urbana-Champaign, Urbana, IL 61801, USA
 ⁵Theoretical Division, Los Alamos National Laboratory, Los Alamos, NM, 87545, USA
 ⁶Center for Theoretical Astrophysics, Los Alamos National Laboratory, Los Alamos, NM, 87545, USA

Submitted to ApJ

ABSTRACT

Neutron star mergers (NSMs) are rapid neutron capture (*r*-process) nucleosynthesis sites that expel matter at high velocities, from 0.1c to as high as 0.6c. Nuclei ejected at these speeds are sufficiently energetic to initiate spallation nuclear reactions with interstellar medium particles. We adopt a thick-target model for the propagation of high-speed heavy nuclei in the interstellar medium, similar to the transport of cosmic rays. We find that spallation may create observable perturbations to NSM isotopic abundances, particularly around the low-mass edges of the *r*-process peaks where neighboring nuclei have very different abundances. The extent to which spallation modifies the final NSM isotopic yields depends on: (1) the ejected abundances, which are determined by the NSM astrophysical conditions and the properties of nuclei far from stability, (2) the ejecta velocity distribution and propagation in interstellar matter, and (3) the spallation cross-sections. Observed solar and stellar *r*-process yields could thus constrain the velocity distribution of ejected neutron star matter, assuming NSMs are the dominant *r*-process source. We suggest avenues for future work, including measurement of relevant cross sections.

Keywords: cosmic rays, *r*-process, nucleosynthesis, nuclear reaction cross sections, nuclear abundances, compact binary stars

Corresponding author: Xilu Wang xwang50@nd.edu, xlwang811@gmail.com

^{*} N3AS Postdoctoral Fellow

1. INTRODUCTION

When particles travel with high speeds ($v \gtrsim 0.1c$; kinetic energy $E_k \gtrsim 5$ MeV) in the interstellar medium (ISM), they are sufficiently energetic to interact with the ISM particles by nuclear reactions. These reactions generally lead to *spallation*: fragmentation processes in which a heavy nucleus emits one or more nucleons, thus reducing its atomic weight.

Nuclear spallation is well studied in the context of cosmic rays. Cosmic rays are highly enriched in Li, Be, and B relative to the ISM or solar system, due to fragmentation of cosmic ray C, N, and O nuclei during propagation in the interstellar medium (e.g., George et al. 2009). The effect of spallation on the cosmic-ray abundance pattern is thus to "fill in the valley" at Li, Be, and B, at the expense of a small reduction in the neighboring CNO peak. Indeed, this effect is not only important for cosmic-ray abundances, but is an important and sometimes dominant nucleosynthesis source for Li, Be, and B generally (Reeves, Fowler & Hoyle 1970; Meneguzzi et al. 1971; Walker, Viola & Mathews 1985; Duncan et al. 1992; Fields et al. 1994; Higdon et al. 1998; Lemoine et al. 1998; Ramaty et al. 2000; Fields et al. 2000; Suzuki & Yoshii 2001).

Spallation may also affect the nucleosynthetic outcomes of individual energetic astrophysical events. For example, spallation reactions can occur in the fast ejecta from supernovae or hypernovae as it interacts with the circumstellar medium (e.g., Fields et al. 2002; Nakamura & Shigeyama 2004). Neutron star mergers (NSMs) are another potential site of interest for spallation studies. Spallation reactions influence the composition of the ultra-heavy cosmic rays (e.g., Binns et al. 2019) that owe their origins to NSMs (Komiya & Shigeyama 2017). They could also alter the abundances of nuclei synthesized in the NSM event, as fast outflowing material from the merger first encounters the ISM. The latter has not yet been considered and is our focus here.

The bulk of heavy nuclei ejected from NSMs are expected to be synthesized through the rapid neutron capture process (r process), which is one dominant nucleosynthesis avenue for heavy elements, especially for those heavier than the iron group (Burbidge et al. 1957; Cameron 1957). In the r process, rapid neutron capture pushes material far from stability and shapes the characteristic abundance pattern with three distinct peaks (at mass numbers $A \sim 80$, $A \sim 130$, and $A \sim 196$), associated with closed shell structures (at neutron numbers N = 50, N = 82, and N = 126). These peaks are clearly seen in the abundance pattern of our solar system (e.g., Lodders 2003; Sneden et al. 2008), where approximately half of the heavy elements have an r-process origin. While the astrophysical site(s) responsible for the galactic tally of r-process elements are still uncertain (see reviews Cowan et al. 1991; Arnould et al. 2007; Cowan et al. 2019; Kajino et al. 2019, and references therein), NSMs are the first verified site of r-process nucleosynthesis (Abbott et al. 2017a,b). The kilonova signal from the multi-messenger event GW170817 indicated lanthanide production from a NSM (Cowperthwaite et al. 2017; Kasen et al. 2017).

In addition, kilonova models of GW170817 suggested that NSMs eject r-process material with high speed that ranges from 0.1c to 0.3c on average (e.g., Kasen et al. 2017; Rosswog et al. 2018; Wollaeger et al. 2018), as expected from previous theoretical work (e.g., Li, &

Paczyński 1998; Tanaka & Hotokezaka 2013). Thus, as this ejecta interacts with interstellar medium, spallation reactions can occur and may influence the overall nucleosynthesis yields from the event. These interactions are the first steps in the NSM ejecta deceleration, and thus are guaranteed to occur for any NSM matter that is eventually stopped and incorporated into observable systems such as stars or the presolar matter.

The r-process ejecta speed and composition are the critical factors for the spallation process, and these depend on the ejecta origin. Simulations find that NSMs permit at least two distinct environments for heavy element production: (1) dynamical ejecta, which is expected to be very neutron-rich (electron fraction $Y_e \sim 0.03-0.2$) and has high speeds in the subrelativistic regime $\sim 0.1-0.3c$, with a velocity tail that can extend to $\sim 0.5-0.6c$ (Bauswein et al. 2013; Hotokezaka et al. 2013; Rosswog et al. 2013; Endrizzi et al. 2016; Lehner et al. 2016; Sekiguchi et al. 2016; Rosswog et al. 2017); (2) a viscous and/or neutrino-driven wind, which is believed to have a range of neutron-richness ($Y_e \sim 0.2-0.5$), but lower velocity ($v \sim 0.1c$) and mass (Chen & Beloborodov 2007; Surman et al. 2008; Dessart et al. 2009; Wanajo et al. 2014; Perego et al. 2014; Martin et al. 2015; Just et al. 2015; Siegel & Metzger 2018). Note that the conditions present in neutrino driven winds are particularly uncertain due to the difficulties in treating neutrinos properly in such environments (e.g., Caballero et al. 2012; Foucart et al. 2015; Malkus et al. 2016; Kyutoku et al. 2018).

In this paper, we investigate the effect of spallation on the shapes of the r-process abundance peaks produced in fast ejecta from a NSM event, and test whether spallation could alleviate the mismatch between simulation results and solar data. In doing so we explore the impact of the many uncertainties in r-process nucleosynthesis. These include nuclear inputs for unstable nuclei, and astrophysical conditions of the merger event (e.g., Mumpower et al. 2016; Kajino & Mathews 2017). These uncertainties write themselves into the r-process abundance patterns as large variations around the second peak ($A \sim 130$) and third peak ($A \sim 195$). For example, the r-process peaks can be narrower and shifted relative to solar data for some astrophysical conditions and choices of input nuclear physics (e.g., Goriely et al. 2011; Korobkin et al. 2012; Just et al. 2015; Rosswog et al. 2017).

The structure of this paper is as follows. The next section describes the methods of our spallation calculation, including the thick-target model for heavy element transport, and the spallation cross-sections of r-process nuclei. Section 3 presents the spallation results for r-process abundances calculated with different astrophysical and nuclear physics inputs. We identify the spallation cross sections with the greatest potential influence on the third peak ($A \sim 195$) region in Section 3.5 and conclude in Section 4.

2. METHOD

To calculate the potential influence of spallation on r-process abundance patterns, we first generate initial abundance patterns of r-process nuclei ejected from a NSM using the nucleosynthesis network code PRISM (see Section 3). We then adopt a thick-target model

for propagation of the r-process nuclei through the ISM, obtaining new abundances of the r-process nuclei after spallation.

2.1. Model Assumptions

We describe the propagation of the r-process nuclei in a one-zone, thick-target "closed-box" model. The initial conditions for our model consists of r-process ejecta from a NSM event. These heavy nuclei propagate into an ISM composed of hydrogen and ${}^4\mathrm{He}$ and experience ionization loss and spallation reactions. The spallation reactions are responsible for the change in the abundance pattern of heavy nuclei as the NSM ejecta is thermalized and eventually incorporated into future generations of stars.

The basic assumptions of our model are as follows:

- 1. The NSM outflows have initial abundances that depend on the adopted r-process nucleosynthesis model and are ejected with a uniform initial speed v.
- 2. The NSM ejecta and ISM are both spatially homogeneous.
- 3. All the *r*-process nuclei will interact with the ISM, i.e., the escape rate is zero. Advection and diffusion loss are also ignored here.
- 4. Among the energy loss mechanisms, ionic/Coulomb losses dominate. For *r*-process nuclei with similar mass number *A* and charge number *Z*, the ionization losses are similar. Thus we treat the energy loss due to ionization as a bulk process that uniformly decelerates the ejecta, and consider spallation as the sole process that will affect the abundance pattern.

We are interested in the early kinematics of the fast heavy particles ejected from a NSM. Similar to supernova remnant evolution, we expect that the ejecta will initially be in free expansion, then sweep enough of the medium that shocks develop and the ejecta decelerates, until finally being stopped (e.g., Chevalier 1977). Our focus is on the very first interactions while the ejecta energies are still high, above the thresholds for spallation. This regime has not been well studied in the NSM case. We select 1 year after the explosion as the starting time for spallation, when the material is in the free expansion phase. At this point *r*-process nucleosynthesis is finished, and the ejecta is expected to be moving with high velocity because there has been little interaction with the medium.

In these early phases during free expansion, we treat the initial particle interactions as scattering events rather than collective hydrodynamic motion. We therefore describe the particle trajectories via the formalism for energy losses of fast particles moving through a medium. Here it is convenient to view the motion in the rest frame of the r-process ejecta, with the medium being an incoming beam of interstellar composition (H and 4 He) moving at speed v_E .

The transport equation for the *r*-process ejecta can be written by adopting the expression for energetic particle propagation used in cosmic-ray studies (e.g., Meneguzzi et al. 1971; Longair 1981; Fields et al. 1994):

$$\partial_t N_E = \partial_E (b_E N_E) + q_E + \text{escape} + \text{advection} + \text{diffusion}$$
 (1)

Here and throughout, E denotes kinetic energy per nucleon, which depends only on the relative velocity between the projectile and target and thus is the same viewed from either frame. The instantaneous number of propagated particles per energy per nucleon at time t is $N_E(E,t)=dN/dE$, thus $N_E\,dE$ is the number of the propagated ejecta nuclei with kinetic energy in the range (E,E+dE). The source function is $q_E=dN/dEdt$ and $b_E=-dE/dt$ is the rate of energy loss (per nucleon). The number flux density is $\phi(E)=v(E)N_E$, where $v(E)=[1-(1+E/(m_pc^2))^{-2}]^{1/2}c$ is the velocity of the ejecta relative to the ISM and m_p is the proton mass, such that for v(E)=0.3c, $E\sim45.29$ MeV.

We simplify the propagation by neglecting the final terms:

$$\partial_t N_E \approx \partial_E (b_E N_E) + q_E \,.$$
 (2)

Here we assume that the only important loss mechanism is the energy loss due to ionization and spallation reactions, and the escape term is thus ignored. This is for the following reasons. (1) In the frame of the NSM ejecta, the incoming ISM particles are the projectiles, and so the energy losses they experience are due to interactions with the ejecta. As discussed in Section 2.1, we are interested at the times after ~ 1 year, when the ejecta is at least partially recombined, so that the energy losses are dominated by ionization losses in a neutral medium. If the ejecta were still fully ionized, Coulomb losses are appropriate. In practice, both of these loss mechanisms are due to the fast particle Coulomb fields, and both share the same scaling with density and particle speed, and have very similar magnitudes (e.g., Mannheim, & Schlickeiser 1994). (2) The pion creation is negligible in the MeV range. (3) Most nuclei in the ejecta are stable or radioactively decay over a much longer timescale than the energy loss timescale of ~ 0.1 Myr (see Appendix B). We omit the advection and diffusion term in our model, as the electro-magnetic interaction time is much shorter than the diffusion and advection times. In addition, spatial uniformity implies that the gradient-driven advection and diffusion terms are zero. The thick-target model presented here neglects secondary particle effects. By assuming that the r-process nuclei lose energy continuously through the propagation, the effect of secondary light nuclei appears only via the elastic scattering energy loss term, and not as a proton/ α particle source term. Secondary heavy particles are also not considered here. While these effects are not large, they would act only to boost spallation, and thus our calculation can be considered a conservative estimate.

We assume the source functions for the projectile/initial r-process nuclei i with mass number A_i , charge Z_i and number abundance Y_i are delta functions in time and in the projectile kinetic energy per nucleon E_0 , i.e., the nuclei in the ejecta are all traveling in the

same speed v_E at time t_0 when ejected by the NSM:

$$q_{i,E}(E,t) = \frac{dN_i}{dt \, dE} = N_{i,0} \, \delta(E - E_0) \, \delta(t - t_0) = N'_{i,0}(E) \, \delta(t - t_0)$$
(3)

where $N_{i,0}$ is the total particle number of r-process element i, and where E_0 is the initial kinetic energy corresponding to initial velocity v.

We calculate a set of spallation reactions $i+j\to\ell+\cdots$ in which projectile i and target j nuclei give rise to products ℓ . A thick-target calculation described in Appendix A gives the product energy spectrum and energy-integrated production rate to be

$$q_{E,ij}^{\ell}(t) = n_j \ \sigma_{ij}^{\ell}(E) \ v(E) \ N_i(E,t).$$
 (4)

$$\frac{dN_{ij}^{\ell}}{dt}(t) = \int q_{E,ij}^{\ell}(t) \ dE \tag{5}$$

where n_j is the ISM number density of targets. Here $\sigma_{ij}^{\ell}(E)$ is the cross section for the production of nuclei ℓ by the reaction between ejecta nuclei i and ISM nuclei j. As discussed in Appendix A and B, since $N_i(E,t) \propto 1/b_i(n_{\rm gas},E)$ (Eq A4), and $b \propto n_{\rm gas}$ (Eqs B9 and B10), the gas density is exactly canceled in the numerator of the Eqs 4 and 5, and thus the final result is *independent of the gas density for the thick-target model*.

Thus the number fraction of the total spallation-produced nuclei ℓ at time t_f to the initial projectile i at time t_0 is:

$$f_i^{\ell} = \sum_{j} f_{i,j}^{\ell} = \sum_{j} \frac{N_{ij}^{\ell}(t_f)}{N_{i,0}} = \sum_{j} y_j \int_{E_x(t_f)}^{E_0} \frac{\sigma_{ij}^{\ell}(E') v(E') dE'}{b_{i,E'}(n_{\text{gas}})/n_{\text{gas}}},$$
 (6)

where E_0 is the initial kinetic energy per nucleon of the projectile nuclei i, which is the maximum kinetic energy of the nuclei. $E_x(t_f)$ is the kinetic energy per nucleon of the projectile nuclei i at time t_f when the nuclei are no longer energetic enough to have spallation reactions. The weighting $y_j = n_j/n_{\rm gas}$ is the fraction by number of ISM particles in the form of $j \in ({\rm H, He})$

Order of magnitude estimate —The fraction $f_{i,j}^{\ell}$ in Eq 6 can also be expressed as an "optical depth" τ_{ij}^{ℓ} :

$$f_{i,j}^{\ell} = \int_{E_x(t_f)}^{E_0} \frac{\sigma_{ij}^{\ell}(E')/m_{\rm u}dE'}{dE'/dX} = \int \kappa_{ij}^{\ell} dX = \tau_{ij}^{\ell}, \tag{7}$$

where the spallation cross section sets in the "opacity" $\kappa_{ij}^{\ell} = \sigma_{ij}^{\ell}/m_{\rm u}$ and $m_{\rm u}$ is the nucleon mass. The path-length of the nucleus i, i.e., the stopping distance or range, is $s_i = \int v dt = \int^E v(E) dE/b_{i,E}$, then $X = \rho_{\rm gas} s = n_{\rm gas} m_{\rm u} s$ is the grammage, which is independent of the medium density and has units ${\rm g/cm^2}$. Our thick target approximation assumes $\tau_{ij}^{\ell} < 1$, and in this limit the optical depth τ_{ij}^{ℓ} or $f_{i,j}^{\ell}$ represents the probability of a spallation reaction occurring, i.e., the fraction of nuclei i that undergo the reaction with the ISM nuclei j.

Now we conduct an order of magnitude calculation to estimate the probability of the reaction between an ISM proton and nucleus i with initial ejecta velocity v. With ionization loss as the dominant channel, we find that $b_{i,E} \sim b_{i,\mathrm{ionic}} \propto Z_i^2 n_{\mathrm{gas}}/(A_i v)$ from Eq B10, and $X_i = \rho_{\mathrm{gas}} \int^E v(E) dE/b_{i,E} \propto A_i v^4/Z_i^2$. We thus have $f_{i,j}^\ell = A_i \sigma_{ij}^\ell/(m_{\mathrm{u}} Z_i^2) X_p$. For v = 0.4c, the grammage for protons traveling through hydrogen is roughly $X_p \sim 1.65~\mathrm{g/cm^2}$. For the third r-process peak, typical cross sections for p+¹⁹⁶Pt are $\sigma_{p,196\mathrm{Pt}}^\ell \sim 1~\mathrm{barn}$ (Kusakabe & Mathews 2018). Therefore $f_{p,196\mathrm{Pt}}^\ell \sim 0.032$, and for each of the dominant channels of spallation reaction p+¹⁹⁶Pt approximately $\sim 3\%$ of ¹⁹⁶Pt is sandblasted. Several such spallation channels will be open, leading to a potential $\sim 10\%$ reduction in ¹⁹⁶Pt and resulting increase in daughter species. This effect is large enough to be of interest, so we proceed with our detailed spallation calculations as described below and show the results in Section 3.

2.3. Spallation Cross Sections of r-Process Nuclei

When propagating through the ISM, the *r*-process nuclei ejected from a NSM lose energy mainly through the ionization of neutral hydrogen (or stopping power) in the MeV range, as discussed in Section 2.2. At the same time, these heavy nuclei also collide with ISM particles and fragment into lighter elements through spallation processes.

In order to understand the production of new nuclei from the initial ejected r-process material interacting with the ISM, we need to know the cross sections for nuclear spallation. Proton spallation reactions have been measured for a few target nuclides (e.g., Hohenberg & Rowe 1970; Brodzinski et al. 1971; Cline & Nieschmidt 1971; Garrett & Turkevich 1973; Perron 1976; Regnier 1979; Fink et al. 1987; Tobin & Karol 1989; Kolsky & Karol 1993; Michel et al. 1997; Yashima et al. 2002; Rejmund et al. 2001; Paradela et al. 2017). However, there are little experimental data available for spallation reactions between a proton or 4 He and a target nuclide which is heavier than iron, in the energy range smaller than $\lesssim 100$ MeV. Therefore in this paper, we adopt the theoretical spallation/inelastic cross sections from TALYS/1.9 (Koning & Rochman 2012, 2019) 1 with default nuclear inputs.

¹ https://tendl.web.psi.ch/tendl 2019/tendl2019.html




Figure 1. Total cross sections for the spallation reaction between ¹⁹⁶Pt and a proton compared to the total ionization loss. The black dotted line is the ionic loss cross section, while the red dotted line is the total non-elastic/spallation cross section calculated using TALYS, the black solid line is the spallation cross section adopted in Kusakabe & Mathews (2018) (KM), and the black dashed line is the spallation cross section adopted in Komiya & Shigeyama (2017) (KS).

Total spallation cross section—The ionization loss dominates over spallation loss, meaning that during propagation, the r-process nuclei mainly lose energy through the electromagnetic interactions. Figure 1 shows the comparisons between the total spallation and ionization loss cross sections for the isotope ¹⁹⁶Pt. The black solid line and dashed line are the spallation cross sections (proton reactions) adopted in Kusakabe & Mathews (2018) (an empirical formula from Letaw et al. (1983)) and Komiya & Shigeyama (2017) (a semiempirical parameterization calculation from SPACS, Schmitt et al. (2014, 2016)), respectively. Compared with TALYS proton results shown in red, these simple formulae give cross sections of same order of magnitude but the differences are not trivial. TALYS contains a variety of options for input nuclear physics such as nuclear level densities, gamma-strength functions, and optical potentials; variations in available inputs result in calculations that differ by at most 10 percent. However, different theory approaches can give larger variations, e.g., NONSMOKER calculations (Bao et al. 2000; Rauscher, & Thielemann 2001; Rauscher 2010)² result in cross section values for the various spallation channels that can differ from TALYS by more than an order of magnitude. For this work we calculate spallation effects using TALYS cross sections (σ_{TALYS}) and with cross sections ten times larger $(10 \times \sigma_{\rm TALYS})$ to roughly account for these uncertainties.

² https://nucastro.org/nonsmoker.html




Figure 2. Individual cross sections for each spallation channel for the reaction between $^{196}\mathrm{Pt}$ and a proton, generated with TALYS. The black solid line is the total spallation cross section, while the colored lines show the cross sections for the individual channels $^{196}\mathrm{Pt}(p,x)A$, where A is the mass number of the final nucleus after spallation. As the projectile energy per nucleon increases, the dominant spallation production channel moves from $A \sim 195$ (dark red circles) to smaller mass numbers, with a wide range of product nuclei at the highest energies.

Cross-sections for each spallation channel—Spallation reactions change a projectile nucleus to a new nucleus with nearby but smaller mass number. Figure 2 shows the cross sections for each spallation channel 196 Pt(p,x). The relevant energy range here is roughly 5-100 MeV; moving from high projectile energy per nucleon to low within this range, the dominant creation channel shifts from producing A=185 to A=195 nuclei. Thus we would expect spallation to shift the r-process abundance pattern peaks to smaller mass numbers, and the results presented in Section 3 confirm this expectation.

2.4. Calculation of the Abundance Change Due to Spallation

From Figure 2, we can see that the number of nucleons that could be removed through spallation reactions depends on the projectile energy per nucleon, or the relative velocity of the projectile and target nuclei. With these spallation cross sections and Eq 6, we calculate the abundance change of the *r*-process nuclei due to spallation, following the number conservation of the total ejecta particles shown in Eq A5.

From our order of magnitude estimate in Section 2.2, we expect spallation to influence any individual abundance Y_i by about 1-10%. Neighboring nuclei tend to have similar spallation cross sections, varying by at most a factor of two. So for regions of the abundance pattern that are fairly flat, any particular mass number A will be depopulated by spallation

roughly at the same rate as it is repopulated by spallation of nuclei with higher A, and any significant rearrangement of the abundance pattern is unlikely. In the regions around the second (110 < A < 140) and third (178 < A < 200) r-process peaks, there are steep abundance changes of an order of magnitude or more. A shift in abundance of 1-10% here can produce a noticeable change to the peak shape, particularly on the lower mass edge. Thus in this work, we focus our spallation calculations exclusively on the two primary peak regions of the main r-process pattern. We expect the spallation effects on other areas of the pattern to be smaller but show similar trends.

For v < 0.1c (corresponding to $E \lesssim 4.73$ MeV/nucleon), the initial kinetic energy is too small to initiate spallation nuclear reactions, so the r-process ejecta needs to be faster than 0.1c for spallation reactions to occur. We consider initial ejecta speeds of v/c = (0.2, 0.3, 0.4, 0.5), corresponding to E = (19, 45, 85, 145) MeV/nucleon). For v = 0.3c, the spallation reactions will remove at most about 5 nucleons from the projectile nuclei (i.e., $A_i - 5 \le A_\ell \le A_i$); for v = 0.4c, the spallation reactions will remove at most 10 nucleons from the projectile nuclei (i.e., $A_i - 10 \le A_\ell \le A_i$). Thus we calculate the spallation effects, i.e., the abundance change of r-process nuclei due to spallation, from the nearby $\sim 10 - 25$ nuclei with heavier mass.

For the third peak (second peak) of the r-process abundance pattern, we check the abundances from A=178 to A=200 (A=110 to A=140), where the abundances peak at $A\sim 196$ ($A\sim 132$) and has a minimum at A=185 (A=110). As the abundances above A=210 (A=150) are negligible compared with the third peak (are smaller and are in a much flatter shape compared with the second peak), we ignore the spallation effects for nuclei with A>210 (150<A<178).

In addition, the ISM is mostly made up of protons ($\sim 75\%$), and α particles (4 He) take up 25% of the ISM total mass, thus the number fraction of protons and 4 He are $y_p \sim 6/7$, $y_\alpha \sim 1/7$. The cross sections for α - 196 Pt spallation reactions show similar trends to p- 196 Pt. Therefore we include both proton and α particle spallation reactions with the initial r-process ejecta nuclei to get the final spallation results.

For each nucleus i interacting with ISM nucleus j through spallation reaction $i+j\to \ell+\cdots$, we calculate $f_{i,j}^\ell$ ($A_\ell=[A_i-n,A_i-0]$). Because the particle number during propagation is conserved (eq. A5), nucleus i produces the same number of nucleus ℓ , thus the loss of nucleus i during the propagation is at the same number as the production of all the nucleus from the spallation reaction of nucleus i, i.e., $f_{i,\text{prop loss}}=\sum_{\ell}(y_pf_{i,p}^\ell+y_\alpha f_{i,\alpha}^\ell)$. The new abundance after spallation is therefore

$$Y_{i,\text{spallation}} = Y_i (1 - f_{i,\text{prop loss}}) + \sum_k (y_p f_{k,p}^i Y_k) + \sum_k (y_\alpha f_{k,\alpha}^i Y_k) . \tag{8}$$

To compare the new abundance pattern with the initial r-process abundance pattern, we compute the spallation abundance change ratio by

$$F_{i,\text{change}} = (Y_{i,\text{spallation}} - Y_i)/Y_i. \tag{9}$$

3. RESULTS

Spallation effects on r-process nuclei ejected from a NSM depend both on the initial r-process nucleosynthesis conditions and on the propagation process. The propagation process is affected by the velocity of the r-process ejecta and spallation cross sections, while the astrophysical conditions and nuclear physics inputs determine the initial r-process abundances generated from nucleosynthesis calculations. Thus both nuclear physics inputs and astrophysical conditions matter for our calculation, and spallation results could place constraints on these conditions in turn.

Nucleosynthesis calculation — In this work, we use the nuclear reaction network code PRISM (Portable Routines for Integrated nucleoSynthesis Modeling) (Mumpower et al. 2016, 2017, 2018) to perform the r-process nucleosynthesis calculations to obtain the abundance patterns for the initial r-process nuclei ejected from a NSM. For our baseline nucleosynthesis calculation set, we begin with the nuclear masses from the 2016 Atomic Mass Evaluation (AME2016) (Wang et al. 2017) if available and FRDM2012 (Möller et al. 2016) otherwise. The Los Alamos National Laboratory (LANL) statistical Hauser-Feshbach code of Kawano et al. (2016) is used to calculate neutron-capture and neutron-induced fission rates for each nuclide. Branchings for β -delayed fission and β -delayed neutron emission are calculated using the LANL QRPA+HF code of Mumpower et al. (2016) as in Möller et al. (2019), with strength functions and half-lives from Möller et al. (2003). Photodissociation is calculated using detailed balance, with one-neutron separation energies calculated directly from the combined FRDM2012 and AME2016 mass dataset. We first explore the case of a symmetric, two-fragment product distribution (symmetric split) for all fissioning nuclei. Finally, we take any values for spontaneous fission and β -decay rates from the Nubase2016 nuclear data evaluation (Audi et al. 2007) to replace any of our theory-based calculations. To gauge the influence of nuclear physics variations on our spallation results, we also adopt β decay rates of Marketin et al. (2016) and neutron capture rates from NON-SMOKER (Bao et al. 2000; Rauscher, & Thielemann 2001; Rauscher 2010)³, in addition to the baseline nuclear reaction rates.

We adopt two kinds of NSM trajectories to compare astrophysical conditions: cold dynamical ejecta (e.g., Goriely et al. 2011; Mumpower et al. 2018) and a low entropy accretion disk wind which is parameterized similar to conditions in McLaughlin & Surman (2005); Surman et al. (2006); Just et al. (2015); Martin et al. (2015); Wanajo et al. (2014); Siegel & Metzger (2018).

3.1. Propagation Parameter Variations

For *r*-process ejecta with a given initial abundance pattern, the spallation abundance change ratio is mainly dependent on the initial ejecta velocity and the spallation cross sections adopted. Here we present our baseline simulation and explore the impact of variations in the initial ejecta velocity and in the spallation cross sections on the final abundances.

³ https://nucastro.org/nonsmoker.html

Baseline spallation calculation—We first consider spallation effects on our baseline r-process abundance pattern using the cold dynamical ejecta conditions described above. Figure 3 shows the resulting abundances of the second and third r-process peaks before and after spallation calculations that assume an initial ejecta velocity of 0.4c, and spallation cross sections from TALYS (σ_{TALYS}) and $10 \times \sigma_{\text{TALYS}}$. We can see that spallation moves the r-process abundance pattern to lower mass numbers, towards the solar data, and smooths the shapes at the left side of the peaks while leaving the right side of the peaks largely unchanged. The spallation effect increases dramatically with an increased spallation cross section.




Figure 3. Abundances in the second peak (110 < A < 140) and third peak (180 < A < 200) of the *r*-process pattern produced with dynamical ejecta of a NSM with initial velocity v = 0.4c and two choices of spallation cross sections, $\sigma_{\rm TALYS}$ (red dashed line) and $10 \times \sigma_{\rm TALYS}$ (orange dotted line). The initial *r*-process abundance pattern from the PRISM simulation is shown in blue and the black points are the solar *r*-process residuals (Arnould et al. 2007). The solar data scales to the ¹⁹⁵Pt abundance from the initial *r*-process simulation.

Third r-process abundance peak—We next explore the effect of ejecta velocity on spallation in the $A\sim 195$ peak region. Figure 4 shows the abundance pattern and the abundance change ratio due to spallation in the third peak region, starting from the same initial cold dynamical ejecta abundance pattern of Figure 3 and considering initial ejecta speeds of 0.3c and 0.5c for the calculation of the spallation effects. Figures 4 and 3 together show that the influence of spallation strongly depends on the velocity of the r-process ejecta, and the abundance pattern changes are non-negligible for ejecta of 0.3c or faster: at 0.3c, spallation brings an $\sim 8\%$ on average of the abundance change and $\sim 60\%$ when the spallation cross section is increased by a factor of 10; at 0.5c, the abundance change can be as high as a factor of 2 for some nuclei. The abundance shape is even flatter than the solar pattern with a velocity of 0.5c and adopted cross sections of $10\times\sigma_{\rm TALYS}$. This suggests that if the actual spallation cross sections are much higher than the predicted TALYS values, the bulk of the ejecta from an r-process event cannot exceed the "speed limit" of about $\sim 0.4-0.5c$.




Figure 4. Spallation effects in the third peak (180 < A < 200) region for the baseline dynamical ejecta simulation assuming initial ejecta velocities of 0.3c (left) and 0.5c (right). Upper panels: Abundances before (blue lines) and after (red/orange lines) spallation, compared to solar data as in Figure 3. Lower panels: The abundance change ratio due to spallation as defined in Eq 9.

Second r-process abundance peak — Figure 5 shows the abundance pattern and the abundance change ratio due to spallation in the second peak region, again for calculations starting from the initial dynamical ejecta abundance pattern of Figure 3 and considering initial ejecta speeds of 0.3c and 0.5c as in Figure 4. The initial abundances include only a main (A>120) r process with no weak/limited r-process component, so the left edge of the second peak is very sharp. Thus the abundance changes due to spallation can be orders of magnitude larger here than for the third peak. Spallation helps to fill the gap with the solar data and smooths the shape of the peak. However, even at the highest initial ejecta speeds where spallation can largely fill in the region to the left of the peak, the fit to solar data remains poor.




Figure 5. Spallation effects in the second peak (110 < A < 140) region for the baseline dynamical ejecta simulation assuming initial ejecta velocities of 0.3c (left) and 0.5c (right). Upper panels: Abundances before (blue lines) and after (red/orange lines) spallation, the solar data scales to the 132 Xe abundance from the initial r-process simulation. Lower panels: The abundance change ratio due to spallation as defined in Eq 9, scaled similarly to Figure 4 to highlight the abundance changes in the second peak itself.

It is important to note that while the third r-process peak is produced only in the most robust neutron-rich environments, a wider variety of conditions can produce the second peak. Thus the region to the left of the second peak is likely filled in with contributions from astrophysical trajectories with higher initial electron fractions Y_e such as from the NSM accretion disk wind (e.g., Holmbeck et al. 2019). While this higher Y_e material may also undergo spallation in the ISM, the effects are likely smaller as the ejecta speeds are expected to be lower. The region to the left of the second peak can also be filled in with the products of heavy fissioning nuclei, an effect we explore in the next section.

3.2. Initial r-Process Abundance Pattern Variations

In Section 3.1, we considered *r*-process ejecta traveling through ISM with different (but still uniform) initial velocities and spallation cross section values, with a fixed initial abundance pattern. Here we repeat the analysis of Section 3.1 with different choices of nuclear physics and astrophysics conditions adopted for the nucleosynthesis, while keeping the ini-

tial velocity of the ejecta at v=0.4c. Both astrophysical conditions and the choice of nuclear data adopted for the nucleosynthesis simulation affect the initial abundance pattern features. This leads to a variance in the potential influence of spallation.




Figure 6. Spallation effects in the third peak (180 < A < 200) region for the baseline dynamical ejecta (left) and hot disk wind (right) simulations assuming initial ejecta velocity of 0.4c. Upper panels: Abundances before (blue lines) and after (red/orange lines) spallation, compared to solar data as in Figure 3. Lower panels: The abundance change ratio due to spallation as defined in Eq 9.

Third r-process abundance peak — Figure 6 compares the abundance patterns and abundance change ratios after spallation for the baseline dynamical ejecta shown in Figure 3 (left) and hot disk wind conditions (right). Hotter r-process freeze-out conditions are characterized by more late-time neutron capture, which produces abundance peaks that can be narrower than and offset from solar data, as shown in the blue lines of Figure 6. Though we recognize that 0.4c may be an unrealistically fast ejecta speed for disk winds, still we consider here whether spallation could possibly alleviate this mismatch. Indeed, with this high initial speed and with the larger $(10 \times \sigma_{\text{TALYS}})$ spallation cross sections, a very narrow, offset initial peak can be smoothed and shifted to produce a reasonable match with solar. In all cases, the effects of spallation are much larger with the sharper peak; the average positive

spallation abundance change is $\sim 200\%$ for the wind example versus $\sim 50\%$ for the cold dynamical ejecta example. Spallation effects are bigger for steeper abundance features.

The r-process proceeds through a region of the nuclear chart where the nuclear properties are highly uncertain (Mumpower et al. 2016). Different choices of nuclear data also yield different initial r-process patterns. We repeat our spallation calculations starting with abundance patterns produced with different choices of nuclear data: β decay rates from Marketin et al. (2016) and neutron capture rates calculated with NONSMOKER (Bao et al. 2000; Rauscher, & Thielemann 2001; Rauscher 2010). Both sets of rates act to broaden the third peak and move the peak position towards solar, resulting in flatter abundance shapes and smaller abundance changes due to spallation compared to those in Figure 6.

Second r-process abundance peak—Figure 7 shows the abundance pattern and abundance change ratio after spallation for the baseline cold dynamical ejecta conditions with two choices of fission yields (Left: simple symmetric split, as in Figure 3; Right: GEF fission yields). The initial r-process abundance data with different fission yields adopted here are from Figure 6 in Vassh et al. (2018), which discusses how fission yields play an important role in shaping the second peak. Compared with the simple symmetric fission yields, GEF (version GEF-2016-V1-2, Schmidt et al. 2016) fission yields fill the huge gap on the left side of the second peak, bringing a flatter abundance pattern and accordingly smaller spallation effect (GEF: $\sim 20\%$ in average; simple symmetric split: $\sim 10000\%$ in average). We also tested Kodama & Takahashi fission yields (Kodama & Takahashi 1975) and found the abundance pattern to be even flatter than solar data, and spallation further enhances the disagreement. Moreover, we tested spallation on the second peak with different astrophysical conditions, including those where fission plays a minor role if any. In general, we find that the effects of spallation on the second peak can vary by $\sim 2-3$ orders of magnitude when different initial conditions are adopted. Thus, the variation in spallation abundance changes due to initial conditions is larger than the variation ($\lesssim 1$ order of magnitude) due to different spallation cross sections. Similar to the third peak, the effects of spallation are the largest where the "cliff" to the left of the peak is sharpest.




Figure 7. Spallation effects in second peak (110 < A < 140) region for the baseline dynamical ejecta simulation assuming initial ejecta velocity of 0.4c, with two choices of fission yields (Left: simple symmetric split; Right: GEF fission yields (Schmidt et al. 2016)). Upper panels: Abundances before (blue lines) and after (red/orange lines) spallation, the solar data scales to the 132 Xe abundance from the initial r-process simulation. Lower panels: The abundance change ratio due to spallation as defined in Eq 9, scaled similarly to Figure 5 to highlight the abundance changes in the second peak itself.

3.3. A Full NSM Simulation

All of our analysis described previously considers only individual astrophysical trajectories with set electron fractions and initial velocities. But in reality, r-process material is ejected from a neutron star merger event with a distribution of velocities, electron fractions, and hydrodynamical conditions. Therefore in this section we adopt the NSM simulation of Bovard et al. (2017) to investigate the influence of spallation on a full NSM event. For our spallation calculations, we use the results of SFHO-M1.35 model (Bovard et al. 2017), which simulates two $1.35M_{\odot}$ neutron stars merging to form a black hole with the ejected mass $M_{ej}=3.53\times 10^{-3}M_{\odot}$. This simulation gives 2253 tracers/trajectories with $\langle v\rangle=0.26c, \langle Y_e\rangle=0.16$. Velocities of the tracers have a large dispersion, ranging from 0.1c to 0.6c. The spallation results on the second and third r-process abundance peaks are shown in Figure 8.




Figure 8. Spallation effects on the second peak (110 < A < 140) and third peak (180 < A < 200) regions of the r-process abundance pattern produced with baseline PRISM calculation for a full NSM simulation Bovard et al. (2017) with two choices of spallation cross sections, $\sigma_{\rm TALYS}$ (red/purple lines) and $10 \times \sigma_{\rm TALYS}$ (orange dotted line). The initial r-process abundance pattern from the PRISM simulation is shown in blue and the black points are the solar r-process residuals (Arnould et al. 2007). The solar data scales to the total abundance of the initial third abundance peak (A = 180 - 200) of the r-process simulation before spallation.

Upper panel: Abundances before (blue lines) and after (red/orange/purple lines) spallation. Red and orange lines are the mass-summed abundance patterns after spallation from each tracer, magenta line is the abundance pattern after spallation with single mass-averaged velocity $v\sim 0.26c$. Lower panel: The abundance change ratio due to spallation as defined in Eq 9, scaled similarly to Figure 4 to highlight the abundance changes in the second and third peaks themselves.

Figure 8 shows the abundance pattern and abundance change ratio after spallation for the NSM simulation we adopted, using the baseline PRISM nucleosynthesis calculations. We add up the initial *r*-process abundances of each tracer based on its mass to get the total initial abundance pattern for the whole merger event shown in the blue solid line. Then we proceed with our spallation calculation in the following two ways for comparison. (1) We perform the spallation calculation on each tracer/trajectory with its associated velocity, then calculate the mass weighted sum using each tracer's abundance pattern after spallation. This gives the total abundance pattern shown in red solid line in Figure 8. We also increase the spallation cross sections by a factor of 10 and repeat the calculation, obtaining the abundance pattern shown in orange dotted line. (2) We perform the spallation calculation on the abundance pattern of the whole combined simulation assuming a mass averaged

initial velocity of 0.26c, similar to the procedure of the previous sections, and obtain the results in purple line.




Figure 9. Spallation effects on the second peak (110 < A < 140) and third peak (180 < A < 200) regions with baseline PRISM calculation for the high speed tracer No.1254 from the NSM simulation with initial ejecta velocity of $v \sim 0.512c$ (Bovard et al. 2017). Upper panels: Abundances before (blue lines) and after (red/orange lines) spallation for the tracer No.1254, compared to solar data as in Figure 8. Lower panels: The abundance change ratio due to spallation as defined in Eq 9, scaled similarly to Figure 8 to highlight the abundance changes in the second and third peaks themselves.

Comparing the red and orange lines with the blue line in Figure 8, we find that spallation produces noticeable changes to the abundance pattern of a full NSM simulation and that the effect increases with an increased spallation cross section. The positive abundance change due to spallation is around 10% on average for the third peak and 100% on average for the second peak. Comparing the red line and purple line, we can see that the estimated effects of spallation on a single combined trajectory are smaller than those of the full calculation. This is mainly due to the contributions of the tracers with the highest speeds, especially for those beyond 0.5c. For example, Figure 9 shows the abundance pattern and abundance change ratio after spallation for tracer No.1254 with v=0.5142c. The blue line is the initial r-process abundance pattern, which has steeper shapes at both the second and third peaks compared to the combined trajectory. The high velocity and the steepness of the abundance peaks together result in larger abundance changes due to spallation. As Figure 8

shows, the faster components of the ejecta do end up influencing the overall yields of a NSM event even though their net mass is small ($\sim 13\%$ of the total ejecta mass).

3.4. Spallation on Elemental Pattern/Elemental Ratio

So far we have focused our attention on the influence of spallation on the isotopic *r*-process pattern. However, isotopic *r*-process abundances are limited to solar system data, while elemental *r*-process abundances are available for a growing number of stars (Sneden et al. 2008; Roederer et al. 2014).




Figure 10. Elemental abundance ratios of Os over Ir (Y(Os)/Y(Ir)) for the dynamical ejecta and hot disk wind simulations assuming initial ejecta velocity of 0.4c (except for the NSM simulation Bovard et al. (2017), which is the mass-summed results of tracers with various velocities), with three choices of adopted nuclear data (PRISM baseline calculation, β decay rates from Marketin et al. (2016) (MKT), and neutron capture rates from NONSMOKER (Bao et al. 2000; Rauscher, & Thielemann 2001; Rauscher 2010)⁵). The ratios are compared with solar data (brown line: solar value, brown region: uncertainty range; Sneden et al. 2008) and stellar observations of the r-process enhanced stars (lines): CS 31081-991 (Hill et al. 2002), HD221170 (Ivans et al. 2006), BD+17°3248 (Cowan et al. 2002), CS 22892-052 (Sneden et al. 2003), HE1523-0901 (Frebel et al. 2007), J0954+5246 (Holmbeck et al. 2018), RAVE J2038-0023 (Placco et al. 2017).

Therefore we also test the effects of spallation on the r-process elemental pattern (abundance Y versus charge number Z) and compare with the stellar observations. For this

analysis, we consider the well-measured third peak elements Os and Ir from r-process enhanced stars, for example, CS 31081-991 (Hill et al. 2002), HD221170 (Ivans et al. 2006), BD+17°3248 (Cowan et al. 2002), CS 22892-052 (Sneden et al. 2003), HE1523-0901 (Frebel et al. 2007), J0954+5246 (Holmbeck et al. 2018), RAVE J2038-0023(Placco et al. 2017). Os/Ir ratios (Y(Os)/Y(Ir)) from solar data and these stellar observations have a wide range (from $\sim 0.8-1.7$). We compare the Os/Ir ratio from different r-process nucleosynthesis simulations (blue circles) to the observation data (lines) in Figure 10. We find that most simulations give the Os/Ir ratio within the solar error bar and consistent with the stellar data. Spallation can adjust this ratio, and we see that although the effect (red stars and triangles) is relatively small, in several cases it moves the Os/Ir ratio more comfortably within the observation range. The one outlier is the case of the cold dynamical ejecta conditions calculated with β decay rates from Marketin et al. (2016), where the simulation ratio is already bigger than the observation range. Therefore for this case, spallation acts in the opposite direction, away from observational data, potentially putting constraints on the ejecta velocity, spallation cross sections, or nuclear data.

3.5. Spallation Cross Section Sensitivity Study

There is little experimental data available for heavy element spallation reactions, especially at the energies smaller than 100 MeV of interest here, and different theoretical estimates can vary by as much as an order of magnitude. As shown in the results throughout this work, the effects of spallation are sensitive to the spallation cross sections adopted. Thus we perform a sensitivity study to identify the nuclei whose spallation cross section adjustments would bring the largest changes to the abundance pattern. We focus on the third peak region, since the second peak shapes are affected by initial nucleosynthesis inputs like fission yields more significantly than spallation cross sections (see Section 3.2).

For our sensitivity study we start with the third peak abundance pattern of the baseline r-process simulation (dynamical ejecta with v=0.4c). We increase the spallation cross section by a factor of 10 for one nucleus at a time, while keeping the other parameters and the cross-sections for all other nuclei unchanged, and repeat the spallation calculation (single-increased scenario). We report the results for the most impactful cross sections in Table 1. Here, we consider the nuclei with increased abundances (i.e., positive change ratios) after spallation, and average their abundance change ratios to obtain the values reported in the table.

We list the top 10 nuclei which result in the largest average positive abundance change ratios. For comparison, Table 1 also shows the average positive abundance change ratios for the calculation with the original TALYS calculated spallation cross section values (original scenario) and the case where TALYS cross sections for all nuclei are increased by a factor of 10 (all-increased scenario). The effects of spallation for the single-increased scenario for 198 Pt, 197 Au, 196 Pt and 195 Pt are $\sim 80\%$ the size of the full effects of the all-increased scenario, suggesting spallation of the third r-process peak is the most sensitive to these four cross sections.

Adopted Cross Sections		Spallation Effect
$\sigma_{ m spallation}$		[percentage]
TALYS value $(1\times)$		23.99
(original)		
10× for all nuclei		138.19
(all-increased)		
$10\times$ for each individual nucleus		
	¹⁹⁸ Pt	114.18
	¹⁹⁷ Au	112.54
	¹⁹⁶ Pt	108.91
	¹⁹⁵ Pt	106.98
	¹⁹⁴ Pt	91.78
	¹⁹³ Ir	64.00
	¹⁹² Os	45.49
	¹⁹¹ Ir	54.67
	¹⁹⁰ Os	46.22
	¹⁸⁹ Os	33.55

Table 1. Spallation cross section sensitivity study starting with the baseline dynamical ejecta simulation with v=0.4c and focused on the third peak of r-process abundance pattern. The table lists the average positive abundance change ratio ("spallation effect") resulting from the original, all-increased, and single-increased scenarios as described in the text. For the single-increases scenario, nuclei with the top 10 most impactful spallation cross sections are shown.

4. DISCUSSIONS AND CONCLUSIONS

In this paper, we report on the construction and results of a thick-target spallation model to test the effects of spallation on the isotopic abundance pattern of nuclei ejected from NSM events. We find that spallation can result in non-negligible changes to relative abundances in the $A\sim 130$ and $A\sim 195$ r-process peak regions of material ejected with speeds of 0.3c and above.

The effects of spallation are to move the abundance pattern towards lower mass numbers and smooth the slope at the left side of the peaks. This effect depends both on the initial r-process nucleosynthesis conditions and on the propagation process. The abundance pattern before spallation is set by the initial astrophysical conditions and the adopted nuclear data; we find that spallation can produce larger changes to abundance peaks that are initially sharper or steeper. Spallation occurs as the ejecta propagates through the ISM, thus faster initial ejecta speeds result in more significant abundance changes. We find that spallation can partially or fully alleviate the mismatch of the second and third r-process abundance peaks compared to solar data. Thus the effects of spallation are non-negligible and need to be considered in addition to other factors for shaping the r-process abundance pattern, especially for the trajectories having steeper abundance peaks and with initial speeds over 0.3c.

Moreover the spallation effects can potentially put constraints on the initial r-process nucleosynthesis conditions and propagation process in turn. The r-process abundance pattern

after spallation should not be flatter than the solar data. Our tests show that such a mismatch can occur for some choices of input nuclear data and for high ejecta speeds, and furthermore suggest that if spallation cross sections are a factor ~ 10 above their theoretical estimates, the bulk r-process ejecta has a 'speed limit' of about 0.4-0.6c, based on its initial abundance shape.

There still remains space to improve our model. A more complete treatment of the particle phase space evolution could take a Boltzmann-like approach to follow the transition from individual particle scattering interactions to the development of shocks. We ignore diffusion here, but in reality, the NSM environment may have strong magnetic fields that could be carried with the ejecta. The motions of both the *r*-process nuclei and the ISM particles may be quite complex if magnetic fields remain strong. In this scenario, the *r*-process nuclei may have more time to undergo spallation reactions, changing the spallation/ionization reaction ratio and thus increasing the spallation effects in a longer timescale. Theoretical work would also benefit from additional studies of spallation reactions and *r*-process nucleosynthesis with different mass models.

As the spallation cross sections we have adopted are from theoretical calculations, our findings call for new experiments. Measurements of spallation reactions of r-process heavy nuclei at relevant energies can pin down the true behavior of the cross sections and branching ratios. Such data will establish the importance of spallation in altering and diagnosing NSM r-process abundances. As seen in Figure 2, the important energy range is $\sim 5-100$ MeV, and the most important projectiles are protons but α particle interactions deserve study as well. The most critical spallation targets are for the $A\sim 196$ nuclides listed in Table 1. Because most spallation reactions occur long after the merger, they involve heavy nuclei at or near stability, and thus sidestep the well-known challenges of studying the highly unstable nuclei essential to the r-process synthesis itself. Thus these measurements could be within reach for appropriate facilities such as FRIB, FAIR and RIKEN .

We are very grateful to Luke Bovard for providing us trajectories from his neutron star merger simulations. We are pleased to thank George Fuller, Wick Haxton, and Shunsaku Horiuchi for the stimulating conversations. This work was supported by U.S. National Science Foundation under grant number PHY-1630782 Focused Research Hub in Theoretical Physics: Network for Neutrinos, Nuclear Astrophysics, and Symmetries (N3AS) (X.W.), and was also supported by the U.S. Department of Energy under Nuclear Theory Contract No. DE-FG02-95-ER40934 (R.S.), DE-AC52-07NA27344 for the topical collaboration Fission In *R*-process Elements (FIRE; N.V. and R.S.), and the SciDAC collaborations TEAMS DE-SC0018232 (T.S., R.S.). This work benefited from conversations stimulated by the National Science Foundation under Grant No. PHY-1430152 (JINA Center for the Evolution of the Elements). M.M. was supported by the US Department of Energy through the Los Alamos National Laboratory. Los Alamos National Laboratory is operated by Triad National Security, LLC, for the National Nuclear Security Administration of U.S. Department of Energy (Contract No. 89233218CNA000001). T.S. was

supported in part by the Los Alamos National Laboratory Center for Space and Earth Science, which is funded by its Laboratory Directed Research and Development program under project number 20180475DR. M.M. was also supported by the Laboratory Directed Research and Development program of Los Alamos National Laboratory under project number 20190021DR.

REFERENCES

- Abbott, B. P., Abbott, R., Abbott, T. D., et al. 2017, Physical Review Letters, 119, 161101
- Abbott, B. P., Abbott, R., Abbott, T. D., et al. 2017, ApJL, 848, L12
- Arnould, M., Goriely, S., & Takahashi, K. 2007, PhR, 450, 97
- Audi, G., Kondev, F. G., Wang, M., Huang,W. J. & Naimi, S. 2017, Chinese Physics C,41, 030001
- Bauswein, A., Goriely, S., & Janka, H.-T. 2013, ApJ, 773, 78
- Bao, Z. Y., Beer, H., Käppeler, F., et al. 2000, Atomic Data and Nuclear Data Tables, 76, 70
- Binns, W., Israel, M. H., Rauch, B. F., et al. 2019, BAAS, 51, 313
- Bovard, L., Martin, D., Guercilena, F., et al. 2017, PhRvD, 96, 124005
- Brodzinski, R. L., Rancitelli, L. A., Cooper, J. A., & Wogman, N. A. 1971, PhRvC, 4, 1250
- Brown, M. D., & Moak, C. D. 1972, PhRvB, 6, 90
- Burbidge, E. M., Burbidge, G. R., Fowler, W. A., & Hoyle, F. 1957, Reviews of Modern Physics, 29, 547
- Caballero, O. L., McLaughlin, G. C., & Surman, R. 2012, ApJ, 745, 170
- Cameron, A. G. W. 1957, The Astronomical Journal, 62, 9
- Chen, W.-X., & Beloborodov, A. M. 2007, ApJ, 657, 383
- Chevalier, R. A. 1977, ARA&A, 15, 175
- Cline, J. E., & Nieschmidt, E. B. 1971, Nuclear Physics A, 169, 437
- Cowan, J. J., Thielemann, F.-K., & Truran, J. W. 1991, PhR, 208, 267
- Cowan, J. J., Sneden, C., Burles, S., et al. 2002, ApJ, 572, 861

- Cowan, J. J., Sneden, C., Lawler, J. E., et al. 2019, arXiv:1901.01410
- Cowperthwaite, P. S., Berger, E., Villar, V. A., et al. 2017, ApJL, 848, L17
- Dessart, L., Ott, C. D., Burrows, A., Rosswog, S., & Livne, E. 2009, ApJ, 690, 1681
- Duncan, D. K., Lambert, D. L., & Lemke, M. 1992, ApJ, 401, 584
- Endrizzi, A., Ciolfi, R., Giacomazzo, B., Kastaun, W., & Kawamura, T. 2016, Classical and Quantum Gravity, 33, 164001
- Fields, B. D., Olive, K. A., & Schramm, D. N. 1994, ApJ, 435, 185.
- Fields, B. D., Olive, K. A., Vangioni-Flam, E., & Cassé, M. 2000, ApJ, 540, 930
- Fields, B. D., Daigne, F., Cassé, M., & Vangioni-Flam, E. 2002, ApJ, 581, 389
- Fink, D., Paul, M., Hollos, G., et al. 1987, Nuclear Instruments and Methods in Physics Research B, 29, 275
- Frebel, A., Christlieb, N., Norris, J. E., et al. 2007, ApJL, 660, L117
- Foucart, F., O'Connor, E., Roberts, L., et al. 2015, PhRvD, 91, 124021
- Garrett, C. K., & Turkevich, A. L. 1973, PhRvC, 8, 594
- George, J. S., Lave, K. A., Wiedenbeck, M. E., et al. 2009, ApJ, 698, 1666
- Goriely, S., Bauswein, A., & Janka, H.-T. 2011, ApJ, 738, L32
- Higdon, J. C., Lingenfelter, R. E., & Ramaty, R. 1998, ApJL, 509, L33
- Hill, V., Plez, B., Cayrel, R., et al. 2002, A&A, 387, 560
- Hohenberg, C. M., & Rowe, M. W. 1970, J. Geophys. Res., 75, 4205
- Holmbeck, E. M., Sprouse, T. M., Mumpower, M. R., et al. 2019, ApJ, 870, 23
- Holmbeck, E. M., Beers, T. C., Roederer, I. U., et al. 2018, ApJL, 859, L24

- Hotokezaka, K., Kiuchi, K., Kyutoku, K., et al. 2013, PhRvD, 87, 024001
- Ivans, I. I., Simmerer, J., Sneden, C., et al. 2006, ApJ, 645, 613
- Just, O., Bauswein, A., Ardevol Pulpillo, R., et al. 2015, MNRAS, 448, 541.
- Kajino, T., & Mathews, G. J. 2017, Reports on Progress in Physics, 80, 084901
- Kajino, T., Aoki, W., Balantekin, A. B., et al. 2019, Progress in Particle and Nuclear Physics, 107, 109
- Kasen, D., Metzger, B., Barnes, J., et al. 2017, Nature, 551, 80.
- Kawano, T., Capote, R., Hilaire, S., et al. 2016, PhRvC, 94, 014612
- Kodama, T., & Takahashi, K. 1975, Nuclear Physics A, 239, 489
- Kolsky, K. L., & Karol, P. J. 1993, PhRvC, 48, 236
- Komiya, Y., & Shigeyama, T. 2017, ApJ, 846, 143.
- Koning, A. J., & Rochman, D. 2012, Nuclear Data Sheets, 113, 2841
- Koning, A. J., & Rochman, D. 2019, Nuclear Data Sheets, 155, 1
- Korobkin, O., Rosswog, S., Arcones, A., et al. 2012, MNRAS, 426, 1940.
- Kusakabe, M., & Mathews, G. J. 2018, ApJ, 854, 183
- Kyutoku, K., Kiuchi, K., Sekiguchi, Y., et al. 2018, PhRvD, 97, 023009
- Lehner, L., Liebling, S. L., Palenzuela, C., et al. 2016, Classical and Quantum Gravity, 33, 184002
- Lemoine, M., Vangioni-Flam, E., & Cassé, M. 1998, ApJ, 499, 735
- Letaw, J. R., Silberberg, R., & Tsao, C. H. 1983, ApJS, 51, 271
- Li, L.-X., & Paczyński, B. 1998, ApJ, 507, L59
- Longair M. S., 1981, High energy Astrophysics, 1st edn., Cambridge Univ. Press, Cambridge and New York, p. 420
- Lodders, K. 2003, ApJ, 591, 1220
- Malkus, A., McLaughlin, G. C., & Surman, R. 2016, PhRvD, 93, 045021
- Mannheim, K., & Schlickeiser, R. 1994, A&A, 286, 983
- Marketin, T., Huther, L., & Martínez-Pinedo, G. 2016, PhRvC, 93, 025805

- Martin, D., Perego, A., Arcones, A., et al. 2015, ApJ, 813, 2
- McLaughlin, G. C., & Surman, R. 2005, NuPhA, 758, 189
- Meneguzzi, M., Audouze, J., & Reeves, H. 1971, A&A, 15, 337
- Michel, R., Bodemann, R., Busemann, H., et al. 1997, Nuclear Instruments and Methods in Physics Research B, 129, 153
- Möller, P., Pfeiffer, B., & Kratz, K.-L. 2003, PhRvC, 67, 055802
- Möller, P., Sierk, A. J., Ichikawa, T., et al. 2016, Atomic Data and Nuclear Data Tables, 109, 1
- Möller, P., Mumpower, M., Kawano, T., & Myers, W. D. 2019, Atomic Data and Nuclear Data Tables, 125, 192
- Mumpower, M. R., Kawano, T., & Möller, P. 2016, PhRvC, 94, 064317
- Mumpower, M. R., Surman, R., McLaughlin, G. C., et al. 2016, Progress in Particle and Nuclear Physics, 86, 86.
- Mumpower, M. R., Kawano, T., Ullmann, J. L., Krtička, M., & Sprouse, T. M. 2017, PhRvC, 96, 024612
- Mumpower, M. R., Kawano, T., Sprouse, T. M., et al. 2018, ApJ, 869, 14
- Nakamura, K., & Shigeyama, T. 2004, ApJ, 610, 888
- Paradela, C., Tassan-Got, L., Benlliure, J., et al. 2017, PhRvC, 95, 044606
- Perego, A., Rosswog, S., Cabezón, R. M., et al. 2014, MNRAS, 443, 3134
- Perron, C. 1976, PhRvC, 14, 1108
- Placco, V. M., Holmbeck, E. M., Frebel, A., et al. 2017, ApJ, 844, 18
- Ramaty, R., Scully, S. T., Lingenfelter, R. E., & Kozlovsky, B. 2000, ApJ, 534, 747
- Rauscher, T., & Thielemann, F.-K. 2001, Atomic Data and Nuclear Data Tables, 79, 47
- Rauscher, T. 2010, PhRvC, 81, 045807 Reeves H., Fowler W. A., Hoyle F., 1970, Natur, 226, 727
- Regnier, S. 1979, PhRvC, 20, 1517
- Rejmund, F., Mustapha, B., Armbruster, P., et al. 2001, NuPhA, 683, 540
- Roederer, I. U., Preston, G. W., Thompson, I. B., et al. 2014, ApJ, 784, 158

Rosswog, S., Piran, T., & Nakar, E. 2013, MNRAS, 430, 2585

- Rosswog, S., Feindt, U., Korobkin, O., et al. 2017, Classical and Quantum Gravity, 34, 104001
- Rosswog, S., Sollerman, J., Feindt, U., et al. 2018, A&A, 615, A132
- Sekiguchi, Y., Kiuchi, K., Kyutoku, K., Shibata, M., & Taniguchi, K. 2016, PhRvD, 93, 124046
- Schlickeiser, R. 2013, Cosmic Ray Astrophysics (Berlin: Springer)
- Schmidt, K.-H., Jurado, B., Amouroux, C., & Schmitt, C. 2016, Nuclear Data Sheets, 131, 107
- Schmitt, C., Schmidt, K.-H., & Kelić-Heil, A. 2014, PhRvC, 90, 064605
- Schmitt, C., Schmidt, K.-H., & Kelić-Heil, A. 2016, PhRvC, 94, 039901
- Siegel, D. M., & Metzger, B. D. 2018, ApJ, 858, 52
- Sneden, C., Cowan, J. J., Lawler, J. E., et al. 2003, ApJ, 591, 936
- Sneden, C., Cowan, J. J., & Gallino, R. 2008, ARA&A, 46, 241

- Surman, R., McLaughlin, G. C., & Hix, W. R. 2006, ApJ, 643, 1057.
- Surman, R., McLaughlin, G. C., Ruffert, M., Janka, H.-T., & Hix, W. R. 2008, ApJL, 679, L117
- Suzuki, T. K., & Yoshii, Y. 2001, ApJ, 549, 303
- Tanaka, M., & Hotokezaka, K. 2013, ApJ, 775, 113
- Tobin, M. J., & Karol, P. J. 1989, PhRvC, 39, 2330
- Vassh, N., Vogt, R., Surman, R., et al. 2018, arXiv:1810.08133
- Walker T. P., Viola V. E., Mathews G. J., 1985, ApJ, 299, 745
- Wanajo, S., Sekiguchi, Y., Nishimura, N., et al. 2014, ApJ, 789, L39
- Wang, M., Audi, G., Kondev, F. G., et al. 2017, Chinese Physics C, 41, 030003
- Wollaeger, R. T., Korobkin, O., Fontes, C. J., et al. 2018, MNRAS, 478, 3298
- Yashima, H., Uwamino, Y., Sugita, H., et al. 2002, PhRvC, 66, 044607

APPENDIX

A. A THICK-TARGET SOLUTION TO THE R-PROCESS EJECTA PROPAGATION

For a thick-target limit, the solution to Eq 2 with the source function Eq 3 gives

$$N_{i}(E,t) = \frac{1}{b_{i}(n_{\text{gas}}, E)} \int_{E}^{\infty} dE' q_{i,E}(E', t')$$

$$= \frac{1}{b_{i}(n_{\text{gas}}, E)} \int_{t}^{\infty} dt' b((n_{\text{gas}}, E'(t')) N'_{i,0}(E'(t')) \delta(t' - t_{0}), \tag{A1}$$

where

$$dt = \frac{dE}{b((n_{\text{gas}}, E))} \Rightarrow dE = b(n_{\text{gas}}, E(t))dt$$
 (A2)

$$\Rightarrow t' = t - \int_{E}^{E'} \frac{dE''}{b(n_{\text{gas}}, E'')}.$$
 (A3)

To do the integral, we need to solve Eq A3 for E when $t'=t_0$, i.e., finding $E=E_x(t)$ so that $\Delta t=t-t_0=\int_{E_x}^{E_0}dE''/b(n_{\rm gas},E'')$. Thus the r-process propagated particles scan down in energy as $\Delta t=t-t_0$ goes from 0 onwards, until reach $E_{\rm threshold}=E_x(t_f)\sim 5$ MeV at time t_f when the particles are no longer energetic enough to have spallation reactions. Therefore we can get $N'_{i,0}(E,t)=N_{i,0}\delta(E-E_x(t))$.

Then we can obtain the spectrum for the propagated r-process nuclei i:

$$N_i(E,t) = \frac{b_i((n_{\text{gas}}, E_x(t))N'_{i,0}(E_x(t))}{b_i(n_{\text{gas}}, E)},$$
(A4)

and the number fraction of the propagated nuclei i at time t_f to the initial projectile i at t_0 is:

$$f_{i,\text{prop}} = \frac{N_{i,\text{prop}}}{N_{i,\text{total}}} = \frac{\int N_i(E, t_f) dE}{N_{i,0}}$$

$$= b_i((n_{\text{gas}}, E_x(t_f)) \frac{1}{b_i(n_{\text{gas}}, E_x(t_f))}$$

$$= 1. \tag{A5}$$

As expected, the particle number is conserved during the propagations.

From the propagated spectrum, we can get the product nuclei ℓ (mass number A_{ℓ} , charge Z_{ℓ}) spectrum (in the lab frame) through the nuclear reactions between projectile nuclei i and nuclei j (mass number A_{j} , charge Z_{j} , number density ratio $y_{j}=n_{j}/n_{\rm gas}$ with n_{j} to be the number density of nucleus j and $n_{\rm gas}$ to be the number density of ISM protons) in the ISM:

$$q_{E,ij}^{\ell}(t) = n_j \sigma_{ij}^{\ell}(E) v(E) N_i(E,t). \tag{A6}$$

$$\frac{dN_{ij}^{\ell}}{dt}(t) = \int q_{E,ij}^{\ell}(t)dE = n_j \int_{E_{-}(t)}^{E_0} dEv(E)\sigma_{ij}^{\ell}(E)N_i(E,t). \tag{A7}$$

Thus the number fraction of the total spallation-produced nuclei ℓ at time t_f to the initial projectile i at time t_0 is:

$$f_{i}^{\ell} = \sum_{j} f_{i,j}^{\ell} = \sum_{j} \frac{N_{ij}^{\ell}(t_{f})}{N_{i,0}} = \sum_{j} \frac{\int_{t_{0}}^{t_{f}} dt dN_{ij}^{\ell}/dt}{N_{i,0}}$$

$$= \sum_{j} \frac{n_{j}}{N_{i,0}} \int_{t_{0}}^{t_{f}} dt \int_{E_{x}(t)}^{E_{0}} dE \frac{b_{i}((n_{\text{gas}}, E_{x}(t))N_{i,0}^{\prime}(E_{x}(t))v(E)\sigma_{ij}^{\ell}(E))}{b_{i}(n_{\text{gas}}, E)}$$

$$= \sum_{j} y_{j} \int_{t_{0}}^{t_{f}} dt v(E_{x}(t))\sigma_{ij}^{\ell}(E_{x}(t))n_{\text{gas}}$$

$$= \sum_{j} y_{j} \int_{E_{x}(t_{f})}^{E_{0}} \frac{\sigma_{ij}^{\ell}(E^{\prime})v(E^{\prime})dE^{\prime}}{b_{i}(n_{\text{gas}}, E^{\prime})/n_{\text{gas}}}$$
(A8)

where E_0 is the kinetic energy per nucleon of the initial projectile nuclei i, which is the maximum kinetic energy of the nuclei. And $\sigma_{ij}^{\ell}(E)$ is the cross section for the production of nuclei ℓ by the reaction between ejecta nuclei i and ISM nuclei j.

B. ENERGY LOSS RATE OF THE R-RROCESS NUCLEI

With the spallation cross sections shown in Section 2.3, we can get the inelastic/spallation energy loss rate self-consistently (assuming the loss is approximated to be continuous):

$$b_{\text{spall}}^{i} = v(E)E\sum_{j} n_{j}\sigma_{ij}^{\text{total}} = n_{\text{gas}}v(E)E\sum_{j} y_{j}\sigma_{ij}^{\text{total}}.$$
 (B9)

Here the cross section $\sigma_{ij}^{\rm total}$ is for all the spallation reactions between projectile nucleus i and target ISM nucleus j. For example, $b_{\rm spall}^{^{196}{\rm Pt}} \sim 2.713 \times 10^{-9}~{\rm MeV/s}$, so the spallation time scale is $\tau_{\rm spall}^{^{196}{\rm Pt}} \sim 196 E/b_{\rm spall}^{^{196}{\rm Pt}} \sim 1.76 {\rm Myr}$, for nuclei $^{196}{\rm Pt}$ traveling with a speed of v=0.3c through the ISM with $n_{\rm gas}=1{\rm cm}^{-3}$, and the spallation mean free path of this nuclei is $\sim 100{\rm kpc}$.

We also include the energy loss rate of *r*-process nuclei due to interactions with ISM (or stopping power; the ionization energy loss is the dominant loss channel) (Schlickeiser 2002):

$$b_{\text{ionic}} \sim 1.82 \times 10^{-13} Z_{\text{eff}}^2 \left(\frac{n_{\text{gas}}}{cm^{-3}}\right)$$

 $\times (1 + 0.0185 \ln(\beta) H(\beta - \beta_0)) \times \frac{2\beta^2}{\beta_0^3 + 2\beta^3} \text{ MeV/As}^{-1}$ (B10)

where $\beta=v/c$, H is the Heaviside step function, $\beta_0=0.01$ is the orbital velocity of electrons in hydrogen atoms. $Z_{\rm eff}=Z(1-1.034\exp(-137\beta Z^{-0.688}))$ is the effective charge (Brown & Moak 1972), which is less than the nucleus's charge Z at low energies. Then the ionization loss time scale $\tau_{\rm ionic}^{196\rm Pt}\sim E/b_{\rm ionic}\sim 0.1$ Myr for the same condition above, and the mean free path of this nuclei is $\sim 10 \rm kpc$. The ionization loss dominate over spallation during the propagation of the r-process ejecta.

Therefore the total energy loss rate per nucleon for nucleus i is $b_i(n_{\rm gas}, E) = b_{\rm spall}^i/A_i + b_{\rm ionic}^i$. Notice that the energy loss rate scales with gas density: $b \propto n_{\rm gas}$ (see Eqs B9 and B10).